

可燃ごみのごみ質調査結果

| 年 度 | | | 令和8年度 | | | |
|-------------------|-----------|---------|-----------|-----------|-------|--|
| 測定年月日 | | | R8. 4. 10 | R8. 5. 11 | 平均 | |
| 天候 | | | 曇 | 晴 | — | |
| 気温[°C] | | | 19.0 | 23.4 | — | |
| ごみの種類組成 (乾ベース) | 紙・布類 | [%] | 56.7 | 50.2 | 53.5 | |
| | 合成樹脂類 | [%] | 16.4 | 20.3 | 18.4 | |
| | ゴム・皮革類 | [%] | 0.0 | 0.0 | 0.0 | |
| | 木・竹・わら類 | [%] | 11.6 | 10.3 | 11.0 | |
| | ちゅう芥類 | [%] | 12.2 | 13.1 | 12.7 | |
| | 不燃物類 | [%] | 1.6 | 3.4 | 2.5 | |
| | その他 | [%] | 1.5 | 2.7 | 2.1 | |
| ごみの種類組成 (湿ベース) | 紙・布類 | [%] | 46.2 | 44.1 | 45.2 | |
| | 合成樹脂類 | [%] | 10.9 | 13.6 | 12.3 | |
| | ゴム・皮革類 | [%] | 0.0 | 0.0 | 0.0 | |
| | 木・竹・わら類 | [%] | 11.8 | 9.2 | 10.5 | |
| | ちゅう芥類 | [%] | 28.9 | 28.9 | 28.9 | |
| | 不燃物類 | [%] | 0.9 | 1.9 | 1.4 | |
| | その他 | [%] | 1.3 | 2.3 | 1.8 | |
| 単位容積重量 | | [kg/m³] | 151 | 142 | 147 | |
| ごみの3成分 | 水分 | [%] | 44.69 | 45.59 | 45.1 | |
| | 灰分(生ごみ) | [%] | 6.33 | 7.27 | 6.8 | |
| | 可燃分 | [%] | 48.98 | 47.14 | 48.1 | |
| 低位発熱量(計算値) | [kcal/kg] | 1,936 | 1,848 | 1,892 | | |
| | [kJ/kg] | 8,104 | 7,736 | 7,920 | | |
| 低位発熱量(実測値) | [kcal/kg] | 2,113 | 2,087 | 2,100 | | |
| | [kJ/kg] | 8,845 | 8,736 | 8,791 | | |
| 灰分(乾燥ごみ) | | [%] | 11.45 | 13.36 | 12.41 | |

可燃ごみのごみ質調査結果

| 年 度 | | | 令和7年度 | | | | | | | | | | | | |
|-------------------|-----------|----------------------|---------|--------|--------|--------|--------|--------|---------|----------|---------|--------|--------|--------|-------|
| 測定年月日 | | | R7.4.11 | R7.5.9 | R7.6.3 | R7.7.4 | R7.8.5 | R7.9.2 | R7.10.3 | R7.11.10 | R8.12.3 | R8.1.7 | R8.2.5 | R8.3.3 | 平均 |
| 天候 | | | 曇 | 晴 | 雨 | 晴 | 晴 | 晴 | 晴 | 晴 | 曇 | 晴 | 晴 | 雨 | — |
| 気温[°C] | | | 19.0 | 21.4 | 21.2 | 30.0 | 33.0 | 31.2 | 26.0 | 19.0 | 15.0 | 8.0 | 10.4 | 13.0 | — |
| ごみの種類組成 (乾ベース) | 紙・布類 | [%] | 45.0 | 51.8 | 49.7 | 55.4 | 53.4 | 56.1 | 48.7 | 55.6 | 52.9 | 51.6 | 59.8 | 54.6 | 52.9 |
| | 合成樹脂類 | [%] | 25.4 | 25.6 | 24.5 | 20.4 | 19.1 | 17.3 | 23.3 | 24.7 | 20.4 | 22.6 | 19.7 | 22.7 | 22.1 |
| | ゴム・皮革類 | [%] | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 2.4 | 0.0 | 0.0 | 0.4 |
| | 木・竹・わら類 | [%] | 13.9 | 10.2 | 8.6 | 12.8 | 11.6 | 13.6 | 14.2 | 8.5 | 12.5 | 10.1 | 5.5 | 7.6 | 10.8 |
| | ちゅう芥類 | [%] | 9.9 | 8.8 | 10.7 | 8.9 | 10.2 | 10.7 | 12.5 | 8.9 | 11.0 | 8.6 | 11.0 | 9.9 | 10.1 |
| | 不燃物類 | [%] | 2.8 | 1.9 | 4.0 | 1.2 | 1.6 | 1.1 | 0.6 | 1.1 | 1.6 | 2.5 | 2.0 | 2.5 | 1.9 |
| | その他 | [%] | 3.0 | 1.7 | 2.5 | 1.3 | 1.8 | 1.2 | 0.7 | 1.2 | 1.6 | 2.2 | 2.0 | 2.7 | 1.8 |
| ごみの種類組成 (湿ベース) | 紙・布類 | [%] | 40.2 | 47.3 | 45.5 | 48.3 | 45.9 | 46.6 | 42.1 | 51.1 | 46.3 | 48.0 | 51.1 | 48.9 | 46.8 |
| | 合成樹脂類 | [%] | 16.9 | 18.1 | 16.3 | 14.1 | 12.7 | 11.7 | 14.7 | 16.1 | 13.8 | 15.9 | 13.7 | 15.9 | 15.0 |
| | ゴム・皮革類 | [%] | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 0.3 |
| | 木・竹・わら類 | [%] | 13.9 | 10.1 | 7.5 | 13.3 | 12.0 | 17.6 | 16.1 | 10.6 | 11.8 | 9.3 | 5.4 | 8.1 | 11.3 |
| | ちゅう芥類 | [%] | 24.9 | 21.7 | 26.4 | 22.2 | 25.2 | 22.3 | 26.2 | 20.5 | 25.5 | 21.4 | 26.7 | 22.9 | 23.8 |
| | 不燃物類 | [%] | 1.5 | 1.1 | 2.2 | 0.7 | 0.9 | 0.6 | 0.3 | 0.6 | 0.9 | 1.4 | 1.1 | 1.4 | 1.1 |
| | その他 | [%] | 2.6 | 1.7 | 2.1 | 1.4 | 1.8 | 1.2 | 0.6 | 1.1 | 1.7 | 2.5 | 2.0 | 2.8 | 1.8 |
| 単位容積重量 | | [kg/m ³] | 155 | 148 | 138 | 148 | 149 | 152 | 138 | 145 | 133 | 126 | 150 | 140 | 144 |
| ごみの3成分 | 水分 | [%] | 46.20 | 42.22 | 44.89 | 41.49 | 44.92 | 45.13 | 47.39 | 46.65 | 43.98 | 42.83 | 43.95 | 44.27 | 44.5 |
| | 灰分(生ごみ) | [%] | 6.28 | 6.36 | 7.14 | 5.85 | 6.08 | 5.59 | 5.02 | 5.43 | 6.38 | 6.87 | 6.61 | 6.89 | 6.2 |
| | 可燃分 | [%] | 47.52 | 51.42 | 47.97 | 52.66 | 49.00 | 49.28 | 47.59 | 47.92 | 49.64 | 50.30 | 49.44 | 48.84 | 49.3 |
| 低位発熱量(計算値) | [kcal/kg] | 1,861 | 2,061 | 1,889 | 2,121 | 1,935 | 1,947 | 1,857 | 1,877 | 1,970 | 2,007 | 1,961 | 1,932 | 1,952 | |
| | [kJ/kg] | 7,790 | 8,627 | 7,907 | 8,879 | 8,100 | 8,150 | 7,773 | 7,857 | 8,246 | 8,401 | 8,209 | 8,087 | 8,169 | |
| 低位発熱量(実測値) | [kcal/kg] | 2,193 | 2,472 | 2,240 | 2,403 | 2,182 | 2,121 | 2,114 | 2,195 | 2,185 | 2,297 | 2,168 | 2,207 | 2,231 | |
| | [kJ/kg] | 9,180 | 10,348 | 9,377 | 10,059 | 9,134 | 8,879 | 8,849 | 9,188 | 9,146 | 9,615 | 9,075 | 9,239 | 9,341 | |
| 灰分(乾燥ごみ) | | [%] | 11.67 | 11.00 | 12.96 | 9.99 | 11.03 | 10.19 | 9.54 | 10.18 | 11.38 | 12.02 | 11.80 | 12.37 | 11.18 |

可燃ごみのごみ質調査結果

| 年 度 | | | 令和6年度 | | | | | | | | | |
|-------------------|-----------|----------------------|-----------|----------|----------|----------|------------|-----------|----------|----------|----------|-------|
| 測定年月日 | | | R6. 4. 18 | R6. 5. 8 | R6. 6. 4 | R6. 7. 9 | R6. 10. 11 | R6. 12. 3 | R7. 1. 8 | R7. 2. 3 | R7. 3. 4 | 平均 |
| 天候 | | | 晴 | 晴 | 曇 | 曇 | 晴 | 晴 | 晴 | 曇 | 曇 | — |
| 気温[°C] | | | 22.0 | 23.0 | 21.2 | 32.5 | 21.0 | 20.5 | 9.8 | 11.0 | 10.6 | — |
| ごみの種類組成 (乾ベース) | 紙・布類 | [%] | 49.00 | 51.8 | 54.5 | 51.7 | 55.6 | 52.4 | 52.3 | 58.2 | 54.6 | 53.3 |
| | 合成樹脂類 | [%] | 22.5 | 25.4 | 19.4 | 22.5 | 19.8 | 24.9 | 20.6 | 17.1 | 22.0 | 21.6 |
| | ゴム・皮革類 | [%] | 2.9 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 0.6 |
| | 木・竹・わら類 | [%] | 6.9 | 6.5 | 9.2 | 12.7 | 10.9 | 9.1 | 8.3 | 11.1 | 9.4 | 9.3 |
| | ちゅう芥類 | [%] | 14.1 | 10.7 | 11.7 | 8.4 | 9.6 | 8.2 | 13.4 | 9.7 | 11.3 | 10.8 |
| | 不燃物類 | [%] | 2.4 | 3.2 | 2.2 | 2.7 | 2.5 | 3.3 | 2.1 | 2.3 | 1.4 | 2.5 |
| | その他 | [%] | 2.2 | 1.1 | 3.0 | 2.0 | 1.6 | 2.1 | 1.7 | 1.6 | 1.3 | 1.8 |
| ごみの種類組成 (湿ベース) | 紙・布類 | [%] | 43.7 | 47.3 | 49.8 | 46.2 | 50.2 | 48.2 | 45.5 | 49.9 | 45.1 | 47.3 |
| | 合成樹脂類 | [%] | 15.2 | 18.2 | 13.2 | 15.8 | 13.7 | 16.3 | 13.3 | 11.6 | 15.7 | 14.8 |
| | ゴム・皮革類 | [%] | 1.7 | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 0.4 |
| | 木・竹・わら類 | [%] | 6.7 | 5.7 | 8.5 | 14.1 | 9.6 | 10.5 | 7.7 | 10.1 | 8.5 | 9.0 |
| | ちゅう芥類 | [%] | 29.5 | 25.00 | 24.7 | 20.2 | 23.3 | 20.7 | 29.8 | 25.3 | 28.4 | 25.2 |
| | 不燃物類 | [%] | 1.3 | 1.9 | 1.2 | 1.6 | 1.4 | 1.8 | 1.1 | 1.3 | 0.8 | 1.4 |
| | その他 | [%] | 1.9 | 1.1 | 2.6 | 2.1 | 1.8 | 2.5 | 1.6 | 1.8 | 1.5 | 1.9 |
| 単位容積重量 | | [kg/m ³] | 149 | 160 | 157 | 151 | 143 | 151 | 146 | 152 | 145 | 150 |
| ごみの成分 | 水分 | [%] | 45.09 | 41.02 | 44.73 | 41.47 | 43.61 | 46.13 | 48.23 | 44.76 | 43.22 | 44.3 |
| | 灰分(生ごみ) | [%] | 6.41 | 6.85 | 6.43 | 7.47 | 6.34 | 6.58 | 6.16 | 6.26 | 6.35 | 6.5 |
| | 可燃分 | [%] | 48.50 | 52.13 | 48.84 | 51.06 | 50.05 | 47.29 | 45.61 | 48.98 | 50.43 | 49.2 |
| 低位発熱量(計算値) | [kcal/kg] | | 1,912 | 2,100 | 1,929 | 2,049 | 1,991 | 1,851 | 1,763 | 1,936 | 2,010 | 1,949 |
| | [kJ/kg] | | 8,004 | 8,791 | 8,075 | 8,577 | 8,334 | 7,748 | 7,380 | 8,104 | 8,414 | 8,159 |
| 低位発熱量(実測値) | [kcal/kg] | | 2,211 | 2,459 | 2,198 | 2,364 | 2,218 | 2,155 | 2,009 | 2,107 | 2,316 | 2,226 |
| | [kJ/kg] | | 9,255 | 10,293 | 9,201 | 9,896 | 9,285 | 9,021 | 8,410 | 8,820 | 9,695 | 9,320 |
| 灰分(乾燥ごみ) | | [%] | 11.68 | 11.62 | 11.64 | 12.76 | 11.25 | 12.21 | 11.89 | 11.34 | 11.18 | 11.73 |

可燃ごみのごみ質調査結果

| 年 度 | | | 令和5年度 | | | | | | | | | | | | |
|-------------------|-----------|----------------------|-----------|-----------|----------|----------|----------|----------|-----------|-----------|-----------|----------|----------|----------|-------|
| 測定年月日 | | | R5. 4. 12 | R5. 5. 11 | R5. 6. 2 | R5. 7. 7 | R5. 8. 4 | R5. 9. 8 | R5. 10. 2 | R5. 11. 7 | R5. 12. 4 | R6. 1. 9 | R6. 2. 2 | R6. 3. 4 | 平均 |
| 天候 | | | 晴 | 曇 | 雨 | 晴 | 晴 | 雨 | 曇 | 晴 | 晴 | 晴 | 曇 | 晴 | — |
| 気温[°C] | | | 20.5 | 21.0 | 23.6 | 30.5 | 29.0 | 29.0 | 26.4 | 24.6 | 16.0 | 9.0 | 11.0 | 10.5 | — |
| ごみの種類組成 (乾ベース) | 紙・布類 | [%] | 50.8 | 47.4 | 46.2 | 52.1 | 49.5 | 50.6 | 53.7 | 61.4 | 55.4 | 52.9 | 57.5 | 50.6 | 52.3 |
| | 合成樹脂類 | [%] | 18.5 | 25.1 | 23.4 | 21.9 | 18.1 | 20.6 | 19.2 | 15.0 | 21.2 | 19.2 | 19.3 | 26.4 | 20.7 |
| | ゴム・皮革類 | [%] | 0.5 | 2.5 | 1.4 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 |
| | 木・竹・わら類 | [%] | 14.1 | 10.9 | 13.0 | 10.6 | 12.6 | 16.1 | 10.8 | 10.1 | 8.2 | 10.5 | 11.1 | 8.6 | 11.4 |
| | ちゅう芥類 | [%] | 12.1 | 10.5 | 10.6 | 10.3 | 13.7 | 9.4 | 11.7 | 9.0 | 9.9 | 12.2 | 8.0 | 11.6 | 10.8 |
| | 不燃物類 | [%] | 2.3 | 2.2 | 3.3 | 4.0 | 2.7 | 1.2 | 1.9 | 2.8 | 2.5 | 3.3 | 2.0 | 1.4 | 2.5 |
| | その他 | [%] | 1.7 | 1.4 | 2.1 | 1.1 | 1.9 | 2.1 | 2.7 | 1.7 | 2.8 | 1.9 | 2.1 | 1.4 | 1.9 |
| ごみの種類組成 (湿ベース) | 紙・布類 | [%] | 46.3 | 45.3 | 41.7 | 48.1 | 42.5 | 47.3 | 49.5 | 53.1 | 48.7 | 46.9 | 50.5 | 46.3 | 47.2 |
| | 合成樹脂類 | [%] | 12.3 | 17.1 | 15.7 | 14.3 | 12.5 | 14.4 | 12.7 | 10.6 | 14.4 | 13.5 | 14.3 | 18.9 | 14.2 |
| | ゴム・皮革類 | [%] | 0.3 | 1.6 | 0.9 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| | 木・竹・わら類 | [%] | 11.9 | 10.4 | 14.0 | 9.8 | 12.1 | 13.2 | 9.5 | 10.8 | 7.3 | 9.1 | 10.7 | 7.2 | 10.5 |
| | ちゅう芥類 | [%] | 26.2 | 22.5 | 23.5 | 24.5 | 28.7 | 22.1 | 24.1 | 22.1 | 25.3 | 26.7 | 21.0 | 25.6 | 24.4 |
| | 不燃物類 | [%] | 1.2 | 1.2 | 1.8 | 2.1 | 1.5 | 0.7 | 1.0 | 1.6 | 1.4 | 1.9 | 1.2 | 0.8 | 1.4 |
| | その他 | [%] | 1.8 | 1.9 | 2.4 | 1.2 | 1.8 | 2.3 | 3.2 | 1.8 | 2.9 | 1.9 | 2.3 | 1.2 | 2.1 |
| 単位容積重量 | | [kg/m ³] | 168 | 166 | 184 | 155 | 167 | 149 | 162 | 165 | 153 | 145 | 129 | 155 | 158 |
| ごみの3成分 | 水分 | [%] | 46.36 | 43.88 | 46.49 | 47.30 | 45.04 | 42.96 | 46.59 | 42.08 | 45.41 | 42.29 | 40.17 | 42.44 | 44.3 |
| | 灰分(生ごみ) | [%] | 6.11 | 6.51 | 6.39 | 7.12 | 6.46 | 6.20 | 6.33 | 6.78 | 6.01 | 7.40 | 6.75 | 6.19 | 6.5 |
| | 可燃分 | [%] | 47.53 | 49.61 | 47.12 | 45.58 | 48.50 | 50.84 | 47.08 | 51.14 | 48.58 | 50.31 | 53.08 | 51.37 | 49.2 |
| 低位発熱量(計算値) | [kcal/kg] | 1,861 | 1,969 | 1,841 | 1,767 | 1,912 | 2,030 | 1,839 | 2,049 | 1,914 | 2,010 | 2,148 | 2,057 | 1,950 | |
| | [kJ/kg] | 7,790 | 8,242 | 7,706 | 7,397 | 8,004 | 8,498 | 7,698 | 8,577 | 8,012 | 8,414 | 8,992 | 8,611 | 8,162 | |
| 低位発熱量(実測値) | [kcal/kg] | 2,099 | 2,377 | 2,175 | 2,090 | 2,120 | 2,316 | 2,106 | 2,219 | 2,162 | 2,276 | 2,414 | 2,487 | 2,237 | |
| | [kJ/kg] | 8,786 | 9,950 | 9,105 | 8,749 | 8,874 | 9,695 | 8,816 | 9,289 | 9,050 | 9,527 | 10,105 | 10,411 | 9,363 | |
| 灰分(乾燥ごみ) | | [%] | 11.40 | 11.60 | 11.95 | 13.52 | 11.76 | 10.87 | 11.86 | 11.71 | 11.01 | 12.83 | 11.28 | 10.76 | 11.71 |